

Allometric relationships of six fish species of the order Characiformes in oxbow lakes on the floodplain of the middle Purus river, Amazon

L. P. DE OLIVEIRA^{1,2,✉}; R. S. DA SILVA²; L. R. VIRGILIO³; F. CORRÊA^{1,2} & L. J. S. VIEIRA^{2,4}

¹ Programa de Pós-Graduação em Ecologia e Conservação, Universidade do Estado de Mato Grosso, Campus de Nova Xavantina. Nova Xavantina, Mato Grosso, Brazil. ² Laboratório de Ictiologia e Ecologia Aquática, Universidade Federal do Acre. Rio Branco, Acre, Brazil. ³ Centro Multidisciplinar, Universidade Federal do Acre. Cruzeiro do Sul, Acre, Brazil. ⁴ Programa de Pós-Graduação em Ecologia e Manejo de Recursos Naturais, Universidade Federal do Acre. Rio Branco, Acre, Brazil.

ABSTRACT. The length-weight relationship (LWR) was estimated for six fish species of the order Characiformes: *Curimatella meyeri*, *Hoplias malabaricus*, *Potamorhina altamazonica*, *Potamorhina latior*, *Pygocentrus nattereri*, *Schizodon fasciatus*, collected from oxbow lakes on the floodplain of the middle Purus River in western Brazilian Amazonia. The specimens were collected in January, May, and September 2012, during both diurnal and nocturnal periods. Allometric coefficients (b) for the studied species ranged between 2.98 and 3.34. The correlation coefficients for the LWR (r^2) ranged between 0.954 and 0.993. The data presented here provide important insights for the development of effective measures for the conservation and management of the study species.

[Keywords: Neotropical region, Amazon basin, Acre, aquatic biodiversity, ichthyofauna]

RESUMO. Relações alométricas de seis espécies de peixes da ordem Characiformes em lagos de ferradura na planície de inundação do rio Purus médio, Amazônia. A relação comprimento-peso (LWR) foi estimada para seis espécies de peixes da ordem Characiformes: *Curimatella meyeri*, *Hoplias malabaricus*, *Potamorhina altamazonica*, *Potamorhina latior*, *Pygocentrus nattereri*, *Schizodon fasciatus*, coletados em lagos de ferradura na planície de inundação do médio rio Purus, no oeste da Amazônia brasileira. As amostras foram coletadas em janeiro, maio e setembro de 2012, durante os períodos diurno e noturno. As espécies apresentaram coeficientes alométricos (b) entre 2.98 e 3.34 e coeficientes de correlação (r^2) variando de 0.954 a 0.993. Os dados aqui apresentados fornecem informações importantes para o desenvolvimento de medidas efetivas para a conservação e manejo das espécies estudadas.

[Palavras chave: região Neotropical, Bacia amazônica, Acre, biodiversidade aquática, ictiofauna]

INTRODUCTION

The Amazon basin has by far the greatest diversity of fishes found in a freshwater system anywhere in the world, with approximately 2406 valid species (Reis et al. 2016; Jézéquel et al. 2020). The Purus River is one of the principal tributaries of the Amazon River, located in the southwestern Amazon basin, where it forms the second largest drainage in the Brazilian state of Acre (SEMA 2012).

Few data have been published on the ichthyofauna of the middle Purus basin, although studies of length-weight relationships (LWR) are available (e.g., Matos et al. 2019; Silva et al. 2020). Estimates of LWR are important tools for the assessment of the growth patterns and nutritional status of fish species (Froese 2006). These estimates can also

provide important input for decision-making on the management and conservation of fish populations. Given these considerations, we investigated the allometric relationships of six species of the order Characiformes: *Curimatella meyeri* (Steindachner 1882), *Hoplias malabaricus* (Bloch 1794), *Potamorhina altamazonica* (Cope 1878), *Potamorhina latior* (Spix and Agassiz 1829), *Pygocentrus nattereri* (Kner 1858), *Schizodon fasciatus* (Spix and Agassiz 1829), found in oxbow lakes on the floodplain of the Purus river, providing important insights into the biology of the study species.

MATERIALS AND METHODS

The source of the Purus River is located in Perú, where it first flows through the departments of Ucayali and Madre de Dios

Order/Family/ Species	N (%)	TL (cm) (min-max)	TW (g) (min-max)	a (95% CL)	b (95% CL)	r ²	P	Growth type	TL (cm) max (Source)	N	Growth type	Published values	
Characiformes													
Anostomidae													
<i>Schizodon fasciatus</i> 115 (16.8) (Spix and Agassiz 1829)	10.0-34.0	15.0-40.7	0.092 (0.0009-0.0941)	2.98 (2.910-3.056)	0.99 <0.01	Isometric	40 (Froese and Pauly 2022)	439	Negative Alometry	45	Negative Alometry	Cella-Ribeiro et al. (2015)	
Curimatidae													
<i>Curimata meyeri</i> 25 (3.6) (Steindachner 1882)	9.5-18.0	14.7-80.0	0.0898 (0.0033-0.0329)	3.01 (2.533-3.373)	0.96 <0.01	Isometric	18 (This study)	45 928	Positive Alometry	45 635	Positive Alometry	Cella-Ribeiro et al. (2015)	
Potamorhina altamazonica (Cope 1878)	8.0-24.0	7.0-145.0	0.0093 (0.0010-0.0727)	2.98 (2.815-3.101)	0.97 <0.01	Isometric	27 (Froese and Pauly 2022)	27	Negative Alometry	27 635	Negative Alometry	Tribuzy-Neto et al. (2015)	
<i>Potamorhina latior</i> 89 (13.0) (Spix and Agassiz 1829)	8.5-22.8	9.0-138.0	0.0048 (0.0034-0.0783)	3.04 (2.832-3.275)	0.95 <0.01	Isometric	65 (Cella-Ribeiro et al. 2015; Froese and Pauly 2022)	65	Negative Alometry	65 67	Negative Alometry	Tribuzy-Neto et al. (2015)	
Erythrinidae													
<i>Hoplias malabaricus</i> (Bloch 1794)	56 (8.2)	4.5-45.0	1.0-1234.0	0.0019 (0.0004-0.0029)	3.17 (3.062-3.240)	0.99 <0.01	Positive Alometry	65 (Froese and Pauly 2022; Giarrizzo et al. 2015)	65	Isometric	65 81	Isometric	Nobile et al. (2015)
Serrasalmidae													
<i>Pygocentrus nattereri</i> (Kner 1858)	303 (42.2)	3.6-28.0	1.02-651	0.0779 (0.0005-0.0898)	3.26 (3.156-3.346)	0.97 <0.01	Positive Alometry	50 (Froese and Pauly 2022)	50	822 Positive Alometry	50 50	Positive Alometry	Cella-Ribeiro et al. (2015); Giarrizzo et al. (2015)

Abbreviations. N (%): number of specimens (numeric percentage of species). TL (min-max): total length (range of values). TW (min-max): total weight (range of values). a: coefficient related to body shape. b: allometric coefficient. CI: confidence interval. r²: correlation coefficient.

Abreviações. N (%): número de espécimes (porcentagem numérica das espécies). CT (min-max): comprimento total (faixa de valores). PT (min-max): peso total (faixa de valores). a: coeficiente relacionado à forma do corpo. b: coeficiente alométrico. CI: intervalo de confiança. r²: coeficiente de correlação.

Table 1. Estimated parameters for the six species of Characiformes collected in oxbow lakes in the floodplain of the middle Purus river, Amazon, Brazil, between January, May and September 2012, and values already published by other authors.

Tabela 1. Parâmetros estimados para as seis espécies de Characiformes coletadas em lagos de ferradura na planície de inundação do médio rio Purus, Amazônia, Brasil, entre Janeiro, Maio e Setembro de 2012, e valores já publicados por outros autores.

before reaching the Brazilian states of Acre, and then, Amazonas, with a total drainage area of 372000 km² (SEMA 2012). The study area (oxbow lakes) is located on the middle Purus between the municipalities of Boca do Acre ($8^{\circ}42'39.75''$ S - $67^{\circ}23'20.40''$ W) and Pauini ($7^{\circ}44'33.32''$ S - $67^{\circ}1'20.35''$ W) in the Brazilian state of Amazonas. The fish specimens were collected from 12 oxbow lakes in January, May and September 2012 (SISBIO #58295-1), using 12 sets of gillnets, 80 m long and 4 m high, with meshes of between 1.5 cm and 12.0 cm. These nets were installed in the early afternoon parallel to the margin of each lake, where they remained for 24 hours, being checked every 4 hours, with the fish being removed, anesthetized with Eugenol (Keene et al. 2008) and later fixed in formalin (10%).

In the laboratory, the species were identified based on taxonomic keys and the voucher specimens (MUFAC-IC 778 until MUFAC-IC 935) were deposited in the ichthyological collection of the Federal University of Acre, in Rio Branco. Each specimen was measured (total length [TL, cm] with an accuracy of 0.01 cm) and weighed (total weight [TW, g], with an accuracy of 0.01 g). The length-weight relationships of the species were determined using the equation $W=a*L^b$ (Le Cren 1951; Froese 2006), where W=total weight of the specimen, L=total length, and a and b are constants. These parameters (a and b) were estimated by linear regression using the formula: $\log(W)=\log(a)+b*\log(L)$, run in PAST 3.0 (Hammer et al. 2001), with the extreme outliers being excluded. The interpretation of b, the slope or growth coefficient, was based on Santos (1978). This parameter allows to determine the type of growth for

each species. The growth type was estimated using the confidence interval (Froese 2006) and the taxonomic classification of the species followed Fricke et al. (2020).

RESULTS

A total of 686 individuals were collected, all representatives of the order Characiformes, including 6 species in 4 families. Half (3) of these species were members of the family Curimatidae, and were represented by 212 individuals, while the other three families were each represented by a single species, that is, the Anostomidae (115 individuals), Erythrinidae (56 fish) and Serrasalmidae (303 individuals). The abundance of each species ranged from 25 individuals (4% of the total) in *Curimata meyeri* to 303 individuals (44%) in *Pygocentrus nattereri*. The largest species collected during the present study was *Hoplias malabaricus*, which had a total length (TL) of up to 45 cm, and the smallest was *P. nattereri*, with a minimum TL of 3.6 cm. The allometric coefficient (b) ranged from 2.98 in *Potamorhina altamazonica* to 3.34 in *Schizodon fasciatus*, while the correlation coefficients for the LWR (r^2) ranged from 0.95 in *Potamorhina latior* to 0.99 in *H. malabaricus* (Table 1, Figure 1).

DISCUSSION

The estimates of the LWR for the six fish species collected from the oxbow lakes on the floodplain of the middle Purus (Table 1) indicated that *S. fasciatus*, *C. meyeri*, *P. altamazonica* and *P. latior* have isometric growth ($b=3$), whereas *H. malabaricus* and *P. nattereri* have positive allometric growth ($b>3$). These values are within a previously defined




Figure 1. Numeric percentage of species of Characiformes collected in oxbow lakes in the floodplain of the middle Purus river, Amazon, Brazil, in January, May and September 2012.

Figura 1. Porcentagem numérica de espécies de Characiformes coletadas em lagoas de ferradura na planície de inundação do médio rio Purus, Amazonas, Brasil, entre janeiro, maio e setembro de 2012.

range for fish species, which range between 2.5 and 3.5, as recorded by Carlander (1969), Froese (2006) and Freitas et al. (2014).

The isometric growth herein recorded for *S. fasciatus* differs from the negative allometric growth recorded for the species in the Amazon River basin (Cella-Ribeiro et al. 2015). In the same study, *C. meyeri* showed positive allometric growth (Cella-Ribeiro et al. 2015), differing from the isometric growth recorded here for this species. Additionally, in contrast to the isometric growth type of our results, *P. altamazonica* and *P. latior* showed negative allometric growth in floodplain lakes in the Amazon basin (Tribuzy-Neto et al. 2015). In *H. malabaricus*, the positive allometric growth was reported in specimens from southern Brazil (Silveira and Vaz-dos-Santos 2015). In contrast, an isometric growth was recorded in *H. malabaricus* from a southeastern Brazilian basin (Nobile et al. 2015) and in a small estuary tributary of the Amazon River basin (Oliveira et al. 2020). In *P. nattereri*, positive allometric growth was also recorded in specimens (Table 1) from rivers in the Amazon basin (Cella-Ribeiro et al. 2015; Giarrizzo et al. 2015).

A range of factors may influence the variation in allometric coefficients, including the ecosystem structure and spatial scale (Froese 2006), given that different environmental conditions (e.g., temperature and land use) result in different life cycle parameters for the same species (Freitas et al. 2017), food

availability (Lima et al. 2017) and nutritional parameters (Freitas et al. 2017) also influence the type of growth of species. Other potentially important factor may be the differences in the techniques used to collect the specimens, since variations in the type of net, mesh or even the sampling schedule may determine more or less subtle variations in the types of specimen or body sizes collected. However, the effective standardization of specimen collection procedures is extremely complicated. Despite these potential limitations, the data presented here provide important insights for the development of effective measures for the conservation and management of the study species, although there is a clear need for further research on the fish fauna of these Neotropical lakes.

ACKNOWLEDGEMENTS. We are grateful to CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior) for financial and technical support, and the SISBIOTA - Top Predators network (CNPq Edital MCT/CNPq/MMA/MEC/CAPES/FNDCT - Transversal Action/FAPs Number 47/2010 - National System of Biodiversity Research - SISBIOTA Brasil). LPO thanks the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for a masters scholarship (financing code: 001). We would also like to thank the research team of the Ichthyology and Aquatic Ecology Laboratory at the Federal University of Acre in Rio Branco for their support.

REFERENCES

- Carlander, K. D. 1969. Handbook of freshwater fishery biology. Vol. 1. Iowa State University Press, Ames, Iowa, United States of America.
- Cella-Ribeiro, A., M. Hauser, L. D. Nogueira, C. R. C. Doria, and G. Torrente-Vilara. 2015. Length-weight relationships of fish from Madeira River, Brazilian Amazon, before the construction of hydropower plants. *Journal of Applied Ichthyology* 31:939-945. <https://doi.org/10.1111/jai.12819>.
- Freitas, T. M. S., J. B. de Sosuza e Souza, B. da S. Prudente, and L. F. A. Montag. 2017. Length-weight relationship in ten fish species from the Nhamundá River, the Amazon Basin, Brazil. *Acta Amazonica* 47:75-78. <https://doi.org/10.1590/1809-4392201601272>.
- Freitas, T. M. S., B. S. Prudente, N. F. Fontoura, and L. F. A. Montag. 2014. Length-weight relationships of dominant fish species from Caxiuã National Forest, Eastern Amazon, Brazil. *Journal of Applied Ichthyology* 30:1081-1083. <https://doi.org/10.1111/jai.12436>.
- Fricke, R., W. N. Eschmeyer, and R. Van Der Laan (eds.). 2020. Eschmeyer's Catalog of Fishes: Genera, Species, References [online]. URL: tinyurl.com/4w3yhxnz.
- Froese, R. 2006. Cube law, condition factor and weight-length relationships: History, meta-analysis and recommendations. *Journal of Applied Ichthyology* 22:241-253. <https://doi.org/10.1111/j.1439-0426.2006.00805.x>.
- Froese, R., and D. Pauly (eds.). 2022. FishBase. World Wide Web electronic publication. URL: fishbase.org.
- Giarrizzo, T., R. R. S. Oliveira, M. C. Andrade, A. P. Gonçalves, T. A. P. Barbosa, A. R. Martins, D. K. Marques, J. L. B. Santos, R. P. S. Frois, T. P. O. Albuquerque, L. F. A. Montag, M. Camargo, and L. Melo de Sousa. 2015. Length-weight and length-length relationships for 135 fish species from the Xingu River (Amazon Basin, Brazil). *Journal of Applied Ichthyology* 31:415-424. <https://doi.org/10.1111/jai.12677>.
- Hammer, Ø., D. A. T. Harper, and P. D. Ryan. 2001. PAST: Paleontological statistics software package for education and data analysis. *Paleontological Statistics*, PAST, 3.0. URL: folk.uio.no/ohammer/past.

- Jézéquel, C., P. A. Tedesco, R. Bigorne, J. A. Maldonado-Ocampo, H. Ortega, M. Hidalgo, K. Martens, G. Torrente-Vilara, J. Zuanon, A. Acosta, E. Agudelo, S. Barrera Maure, D. A. Bastos, J. Bogotá Gregory, F. G. Cabeceira, A. L. C. Canto, F. M. Carvajal-Vallejos, L. N. Carvalho, A. Cella-Ribeiro, R. Covain, C. Donascimiento, C. R. C. Dória, C. Duarte, E. J. G. Ferreira, A. V. V. Galuch, T. Giarrizzo, R. P. Leitão, J. G. Lundberg, M. Maldonado, J. I. Mojica, L. F. A. Montag, W. M. Ohara, T. H. S. Pires, M. Pouilly, S. Prada-Pedreros, L. J. de Queiroz, L. Rapp Py-Daniel, F. R. V. Ribeiro, R. Ríos Herrera, J. Sarmiento, L. M. Sousa, L. F. Stegmann, J. Valdiviezo-Rivera, F. Villa, T. Yunoki, and T. Oberdorff. 2020. A database of freshwater fish species of the Amazon Basin. Scientific Data 7:1-9. <https://doi.org/10.1038/s41597-020-0436-4>.
- Keene, J. L., D. L. G. Noakes, R. D. Moccia, and C. G. Soto. 2008. The efficacy of clove oil as an anaesthetic for rainbow trout, *Oncorhynchus mykiss* (Walbaum). Aquaculture Research 29:89-101. <https://doi.org/10.1046/j.1365-2109.1998.00927.x>.
- Le Cren, E. 1951. The Length-Weight Relationship and Seasonal Cycle in Gonad Weight and Condition in the Perch (*Perca fluviatilis*). British Ecological Society 20:201-219. <http://doi.org/10.2307/1540>.
- Lima, E. M. M., K. N. S. Sousa, P. R. B. Santos, L. A. R. Ferreira, A. F. Rodrigues, and A. dos S. Pantoja. 2017. Relação peso-comprimento e fator de condição da pescada branca (*Plagioscion squamosissimus*, Heckel 1840) comercializada no município de Santarém, Pará, Brasil; Biota Amazônia 7:44-48.
- Matos, O. F., D. V. Pereira, J. Aguiar-Santos, A. da S. Sampaio, C. E. de Carvalho Freitas, and F. K. Siqueira-Souza. 2019. Length-weight relationships of five fish species from lakes of the Central Amazonian floodplains. Journal of Applied Ichthyology 35:799-801. <https://doi.org/10.1111/jai.13901>.
- Nobile, A. B., E. M. Brambilla, F. P. de Lima, D. Freitas-Souza, I. L. Bayona-Perez, and E. D. Carvalho. 2015. Length-weight relationship of 37 fish species from the Taquari River (Paranapanema Basin, Brazil). Journal of Applied Ichthyology 31:580-582. <https://doi.org/10.1111/jai.12761>.
- Oliveira, M. S. B., L. M. A. Silva, L. Prestes, and M. Tavares-Dias. 2020. Length-weight relationship and condition factor for twelve fish species from the Igapó Fortaleza basin, a small tributary of the Amazonas river estuary. Acta Amazonica 50:8-11. <http://doi.org/10.1590/1809-4392201900702>.
- Reis, R. E., J. S. Albert, F. Di Dario, M. M. Mincarone, P. Petry, and L. A. Rocha. 2016. Fish biodiversity and conservation in South America. Journal of Fish Biology 89:12-47. <https://doi.org/10.1111/jfb.13016>.
- Santos, E. P. 1978. Dinâmica de populações aplicada à pesca e piscicultura. São Paulo, Edusp. 78.
- SEMA. 2012. Plano estadual de recursos hídricos do Acre. Page (SEMA, Ed.). First edition. Rio Branco.
- Silva, R. S., L. R. Virgilio, F. Corrêa, and L. J. S. Vieira. 2020. Length-weight relationships of fish species from oxbow lakes on the floodplain of the middle Purus River in western Brazilian Amazon. Journal of Applied Ichthyology 36: 256-258. <https://doi.org/10.1111/jai.14013>.
- Silveira, E. L., and A. M. Vaz-dos-Santos. 2015. Length-weight relationships for 22 neotropical freshwater fishes from a subtropical river basin. Journal of Applied Ichthyology 31:552-554. <https://doi.org/10.1111/jai.12699>.
- Tribuzy-Neto, I., & K. G. Conceição, F. K. Siqueira-Souza, and C. E. Freitas. 2015. Length-Weight Relationship of Eleven Fish Species of the Amazonian Floodplain Lakes. Revista Colombiana De Ciencia Animal 7:77-79. <https://doi.org/10.24188/recia.v7.n1.2015.425>.